
Modification of the composition and density of Mercury from late accretion



Ryuki Hyodo

JAXA

Today's Contents

- Introduction

What is "late accretion"?

Late accretion depends on planet formation model

- Methods → Consequences of late accretion


How significantly could Mercury's surface be eroded? How much of impactors' materials could be implanted on Mercury's surface? What are the fate of impactors? — Melted? Vaporized?

- Summary

- Discussion


What should we do in the future?

What is "late accretion"?



Late accretion leads to numerous crater-forming impacts: that is, a *statistical* approach can be applied.

*A giant impact is a *stochastic* process




Late accretion depends on planet formation model

Model	Total mass of impactors to Mercury	Impact velocity		
Decline model				
Time: $4.5 - 4.0$ Ga	$3.3^{+4.6}_{-2.6} \times 10^{20}$ kg (Brasser et al., 2020)	Equation 1 (< $v_{\rm imp}$ >= 36 km $^{-1}$)		
Classical LHB		•		
Time: 3.95 – 3.85Ga	0.3×10^{20} kg (Abramov et al., 2013)	$33 \text{ km s}^{-1} \text{ or } 43 \text{ km s}^{-1} \text{ (Mojzsis et al., 2018)}$		
Sawtooth LHB				
Time: 4.1 – 3.7Ga	0.084×10^{20} kg (Abramov and Mojzsis, 2016)	$33 \text{ km s}^{-1} \text{ or } 43 \text{ km s}^{-1} \text{ (Mojzsis et al., 2018)}$		

 $M_{\text{imp,tot}}=8\times10^{18}\sim8\times10^{20}\text{ kg}$

 $\langle v_{\rm imp} \rangle = 30 \sim 40 \text{ km/s}$



Total mass of impactors & typical impact velocity depend on planet formation models.

A strategic consideration

Every few years(!), models are updated.. Total mass of impactors ($M_{\text{tot,imp}}$) & typical impact velocity (v_{imp}) could be changed.


Therefore, instead of studying a *specific* case, I try to derive scaling laws for the *consequences* of late accretion as a function of $M_{\text{tot,imp}}$ and v_{imp} .

Here, the studied consequences are

- (1) How significantly could Mercury's surface be eroded?
- (2) How much of impactors' materials could be implanted on Mercury's surface?
- (3) What are the fate of impactors? Melted? Vaporized?

Preparation

Hyodo & Genda (2020, 2021), ApJ



Scaling laws

Erosion mass from target

$$\frac{M_{\text{HG20,esc,imp}}(>v_{\text{esc}})}{m_{\text{imp}}} = C_{\text{HG20,imp}}(\theta) \left(\frac{v_{\text{esc}}}{v_{\text{imp}} \sin(\theta)}\right)^{-3\mu_{\text{HG20,imp}}(\theta)}$$

Accretion mass from impactor

$$\begin{split} \frac{M_{\text{HG20,acc,imp}}(< v_{\text{esc}})}{m_{\text{imp}}} \\ &= 1 - C_{\text{HG20,imp}}(\theta) \left(\frac{v_{\text{esc}}}{v_{\text{imp}} \sin \theta}\right)^{-3\mu_{\text{HG20,imp}}(\theta)} \\ *C, \mu: \text{coefficients} \end{split}$$

 v_{imp} : impact velocity, m_{imp} : impactor mass, v_{esc} : escape velocity

Scaling laws are derived for erosion and accretion by cratering impacts

Monte-Carlo Statistical Model of Late Accretion

Erosion mass from target

$$\frac{M_{\text{HG20,esc,imp}}(>v_{\text{esc}})}{m_{\text{imp}}} = C_{\text{HG20,imp}}(\theta) \left(\frac{v_{\text{esc}}}{v_{\text{imp}} \sin(\theta)}\right)^{-3\mu_{\text{HG20,imp}}(\theta)}$$

Accretion mass from impactor

$$\frac{M_{\text{HG20,acc,imp}}(< v_{\text{esc}})}{m_{\text{imp}}}$$

$$= 1 - C_{\text{HG20,imp}}(\theta) \left(\frac{v_{\text{esc}}}{v_{\text{imp}} \sin \theta}\right)^{-3\mu_{\text{HG20,imp}}(\theta)}$$

Late accretion — inevitable

(e.g., $M_{\text{imp,tot}}=8\times10^{18}~8\times10^{20}$ kg, $v_{\text{imp}}=30~40$ km/s)


*C, μ : coefficient

What is the consequences of late accretion?

- (1) How significantly could Mercury's surface be eroded?
- (2) How much of impactors' materials were implanted on Mercury's surface?
- (3) What are the fate of impactors? Melted? Vaporized?

We will derive scaling laws as a function of Mimp,tot & Vimp

(so that any planet formation models can be applied)




Consequence I:


How significantly could Mercury's surface be eroded?

- Can late accretion significantly change the core fraction?
- What is the total crater are?

Can late accretion significantly change core fraction?

Figure: Total impactors mass necessary to change the core fraction from $f_{\text{ini,core}}$ to f_{core} =0.7







Late accretion itself *cannot* significantly change the core fraction.

Can late accretion significantly change core fraction?

Figure: Total impactors mass necessary to change the core fraction from $f_{\text{ini,core}}$ to f_{core} =0.7





However, the *surface/crust* could be significantly eroded.

Derived scaling law

Depth of crustal erosion by late accretion (D_{ero}):

$$D_{\text{ero}} \simeq \frac{M_{\text{esc,tot}}}{4\pi\rho R_{\text{Mer}}^2} \simeq 1 \text{km} \times \left(\frac{v_{\text{imp}}}{36 \text{km/s}}\right)^{2.2} \left(\frac{M_{\text{imp,tot}}}{10^{20} \text{kg}}\right)$$


From the existing models

*assuming today's Mercury size

$$M_{\text{imp,tot}}=8\times10^{18}\sim8\times10^{20} \text{ kg & } v_{\text{imp}}=30\sim40 \text{ km/s}$$

 $\rightarrow 50 \text{m}\sim50 \text{ km surface was eroded}$

What is the total crater are?




Derived scaling law

Total crater area to Mercury's surface area (i.e., the degree of resurfacing)

$$\frac{S_{\text{tot,cra}}}{S_{\text{Mer}}} = a \times \left(\frac{M_{\text{imp,tot}}}{10^{20} \text{kg}}\right)^b$$

a & b: coefficients from pi-scaling law and Monte-Carlo simulations (see Hyodo+2021)



From the existing models

If
$$M_{\text{imp,tot}} > 4 \times 10^{19} \sim 5 \times 10^{19} \text{ kg}$$

 $\rightarrow \text{global resurfacing}$
(i.e., $S_{\text{tot,cra}}/S_{\text{Mer}} > 1$)

The oldest surface age for Mercury:

$$\sim 4.0 - 4.1$$
 Ga


(Marchi et al., 2013; Werner, 2014)

Consequence II:

How much of impactors' materials were implanted on Mercury's surface?

*This is the process to deliver exogenic impactor's materials, including the volatile-bearing materials, onto Mercury's surface.

How much of impactors' materials were implanted on Mercury's surface?




Derived scaling law

If $M_{\text{imp,tot}}$ is impacted, $M_{\text{acc,imp,tot}}$ is implanted onto Mercury:

$$M_{\text{acc,imp,tot}} = \gamma M_{\text{imp,tot}}$$

 γ =0.39-0.48 for v_{imp} =30-40 km/s




Consequence III:

What are the fate of impactors? Melted? Vaporized?

*This is the very first step to understand potential geochemical features produced by late accretion impacts.

What are the fate of impactors? — Melted? Vaporized?



Using the lever rule, the melt/vapor fraction estimated

- Entropy calculation along Hugoniot
- Basalt assumed with the initial temperature 293K



Used impact conditions

- 30~40 km/s with 45-degree impact
- Vertical component is used (Pierazzo and Melosh, 2000)



~50% of the impactor materials are vaporized; rest melted *This is a cratering impact; thus a local phenomenon


Summary

Hyodo, Genda & Brasser (2021) Icarus

Existing models

 $M_{\rm imp,tot} = 8 \times 10^{18} - 8 \times 10^{20} \text{ kg}$

 $v_{\rm imp} = 30\sim 40 \text{ km/s}$



How significantly could Mercury's surface be eroded?

$$D_{\text{ero}} \simeq \frac{M_{\text{esc,tot}}}{4\pi\rho R_{\text{Mer}}^2} \simeq 1 \text{km} \times \left(\frac{v_{\text{imp}}}{36 \text{km/s}}\right)^{2.2} \left(\frac{M_{\text{imp,tot}}}{10^{20} \text{kg}}\right)$$

→ 50m~10 km of crust removed

How much impactors materials were implanted?

$$M_{\rm acc,imp,tot} = \gamma M_{\rm imp,tot}$$


→ >3×10¹⁸ kg exogenic materials implanted

What is the total crater area to Mercury's surface area?

$$\frac{S_{\text{tot,cra}}}{S_{\text{Mer}}} = a \times \left(\frac{M_{\text{imp,tot}}}{10^{20} \text{kg}}\right)^b$$

 \rightarrow M_{imp,tot} > 5 × 10¹⁹ kg leads to global resurfacing

What should we do in the future?



Can we find an evidence of late accretion?

- Need to study a local geochemical/geophysical evolution of impact melt/vapor after the impact togethr with observational evidences to constrain the impactor's composition (e.g., icy or rocky impactor).
- Understanding late accretion leads to a better understanding of the solar system formation.

Today's understanding

Study of the Solar system formation

Better constrain

— model update —

Future observation

Surface observation by BepiColombo

Thisstudy

Late accretion onto Mercury

- i.e., total mass, impact velocity-

Thisstudy

What are the consequences of late accretion?

- implantation? erosion? vaporization? -

→ We derived scaling laws (a function of M_{tot,imp} & V_{imp})

Detail modeling of each impact is requred.
— e.g., impact melt/vapor geochemistry —



Future study

What is the consequences of a big impact?

(a stochastic event)

Mercury is Special among Terrestrial Planets

Hyodo & Genda (2021) *ApJ*

Name (rocky/icy)	$v_{\rm imp} [{\rm km \ s}^{-1}]$ $(v_{\rm imp} [v_{\rm esc}])$	Description (Reference) Instability phase (e.g., Hyodo et al. 2021)	
Mercury (rocky)	$\sim 36 \text{ km s}^{-1}$ ($\sim 8.5 v_{\text{esc}}$; $v_{\text{esc}} \simeq 4.25 \text{ km s}^{-1}$)		
Venus (rocky)	$\sim 26 \text{ km s}^{-1}$ ($\sim 2.5 v_{\rm esc}$; $v_{\rm esc} \simeq 10.4 \text{ km s}^{-1}$)	Instability phase (e.g., Mojzsis et al. 2019)	
Earth (rocky)	$\sim 22 \text{ km s}^{-1}$ ($\sim 2.0 v_{\text{esc}}$; $v_{\text{esc}} \simeq 11.2 \text{ km s}^{-1}$)	Instability phase (e.g., Mojzsis et al. 2019)	
Moon (rocky)	$\sim 19 \text{ km s}^{-1}$ ($\sim 8.0 v_{\rm esc}$; $v_{\rm esc} \simeq 2.4 \text{ km s}^{-1}$)	Instability phase (e.g., Brasser et al. 2020)	
Mars (rocky)	$\sim 13 \text{ km s}^{-1}$ ($\sim 2.6 v_{\rm esc}$; $v_{\rm esc} \simeq 5.0 \text{ km s}^{-1}$)	Instability phase (e.g., Mojzsis et al. 2019	
Ceres (rocky)	$\sim 5 \text{ km s}^{-1}$ ($\sim 9.8 v_{\rm esc}$; $v_{\rm esc} \simeq 0.5 \text{ km s}^{-1}$)	Today's population (e.g., Bottke et al. 1994)	
Vesta (rocky)	$\sim 5 \text{ km s}^{-1}$ ($\sim 13.9 v_{\rm esc}$; $v_{\rm esc} \simeq 0.35 \text{ km s}^{-1}$)	Today's population (e.g., Bottke et al. 1994)	
Rhea (icy)	$\sim 15 \text{ km s}^{-1}$ ($\sim 23.6 v_{\rm esc}$; $v_{\rm esc} \simeq 0.64 \text{ km s}^{-1}$)	Instability phase (e.g., Wong et al. 2020)	
Europa (icy)	$\sim 24 \text{ km s}^{-1}$ ($\sim 12.0 v_{\rm esc}$; $v_{\rm esc} \simeq 2.0 \text{ km s}^{-1}$)	Instability phase (e.g., Wong et al. 2020)	
Pluto (icy)	$\sim 2 \text{ km s}^{-1}$ ($\sim 1.7 v_{\rm esc}$; $v_{\rm esc} \simeq 1.2 \text{ km s}^{-1}$)	Today's population (e.g., Dell'Oro et al. 2013	

During late accretion, only Mercury experiences a net erosive outcome.

